


The Case for a Uranus Orbiter and How it Addresses Satellite Science

Mark Hofstadter

Jet Propulsion Laboratory/California Institute of
Technology

Report to the Decadal Survey Satellites Panel,
24 August 2009



Near-IR image from VLT ANTU, © ESO

JPL Contributors

This talk is based on an internal JPL study of missions to Uranus. Co-authors are listed below.

A White Paper, “The Case for a Uranus Orbiter,” is currently being circulated.

Science Co-I's:

Kevin Baines
Shawn Brooks
Leigh Fletcher
A. James Friedson
Robert Lock
Neil Murphy
Glenn Orton
Robert Pappalardo
Nicole Rappaport
Christophe Sotin
Daniel Wenkert


Rapid Mission Architecture Team:

Tom Spilker (Study Lead)
William Smythe
Robert Moeller
Chester Borden
Erick Sturm
Robert Miyake
Paul Stella
Robert Kinsey
Chuck Baker

This research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

Outline of Today's Talk

- I) Introduction
- II) The scientific importance of ice giants
- III) General results from our study of Uranus missions
- IV) Specifics of a New Frontiers-class Uranus Interior Mission
- V) Conclusions



Outline of Today's Talk


I) Introduction

II) The scientific importance of ice giants

III) General results from our study of Uranus missions

IV) Specifics of a Uranus Interior Mission

V) Conclusions




Ice Giant Science Questions (Page 1 of 4)

What is the bulk composition and interior structure of the ice giants?

This is the most important question to address, as it defines what an ice giant is. It influences our understanding of

- The proto-planetary nebula (composition and dynamics),
- Planetary formation (how and where planets form),
- Thermal and chemical evolution of planets (heat flow, interior convection),
- Extra-solar planets.



Images © C.J. Hamilton

Ice Giant Science Questions (Page 2 of 4)

Where and how is the magnetic field generated?

The magnetic field is important for understanding

- Upper atmospheric composition and energy balance,
- The interior structure (conductive and convective regions),
- The dynamo generation process.




Image courtesy F. Bagenal

Ice Giant Science Questions (Page 3 of 4)

What is the nature of internal heat transport within Uranus?

Uranus is emitting essentially no internal heat, perhaps due to density variations inhibiting convection. This impacts Uranus'

- Evolution,
- Interior structure and circulation,
- Atmospheric dynamics and composition,
- The dynamo generation process.

Internal Heat	Jupiter	Saturn	Uranus	Neptune
$W / m^2 \times 10^{11}$	5440 ± 430	2010 ± 140	42 ± 47	433 ± 46
$W / kg \times 10^{10}$	1.7	1.5	0.04	0.32
Internal / Absorbed Solar	0.7	0.8	0.08	1.6

Based on Guillot 2005



Outer Planet Satellites

Jupiter
Saturn
Uranus
Neptune



Europa

Ganymede

Callisto



Earth

H_2O, CO_2



Enceladus

Titan

NH_3, CO, CH_4



Triton

N_2



Ice Giant Science Questions (Page 4 of 4)

What is the nature of the ice-giant satellites?

- Why is their size distribution much different than those of the gas giants?
- Is their composition and structure distinct from those of the gas giants?
- What geologic processes account for the relatively young and tortured (e.g. Miranda) surfaces?
- Are liquid oceans present? (Note that the magnetic field orientation of ice giants optimizes detection of induced currents in the satellites.)

Why a Mission to an Ice Giant?

In the parameter space of “all possible planets,” Uranus and Neptune occupy a region we know very little about. They have an important story to tell about planetary formation and evolution. Learning about them is particularly important if we are to understand extra-solar planetary systems.

Either ice giant can serve as a model for this poorly-understood class of planet, and both have unique features worthy of study.

All major categories of objects in our solar system have a dedicated mission currently flying, except for the ice giants.

A New Frontiers (or Small Flagship) mission launched to an ice giant in the next decade is the only way to dramatically advance our understanding of these objects in our professional lifetimes.

Why Uranus Instead of Neptune?

- Uranus' interior structure and internal heat flow are most challenging to our understanding of planetary formation and evolution, and better constrain our models.
- Uranus is closer, allowing for
 - Shorter flight times (reduced cost and greater reliability),
 - More sunlight (for imaging and power),
 - Better ground-based supporting observations.
- The uranian satellite system may be our solar system's only surviving example of an ice giant system.
- Uranus' atmosphere and satellites experience extreme seasonal forcing due to the system's 98° axial tilt.
- It allows a scientifically compelling mission in our lifetimes.

Outline of Today's Talk


I) Introduction

II) The scientific importance of ice giants

III) General results from our study of Uranus missions

IV) Specifics of a Uranus Interior Mission

V) Conclusions




The JPL Study

Late last summer, we undertook a small study to explore the feasibility of a solar powered mission to Uranus. Many results are relevant to nuclear powered missions as well.

Our primary focus was on New Frontiers, but we also considered higher-cost options.

We engaged JPL's "Rapid Mission Architecture" (RMA) process, which compliments the more familiar Team-X studies. RMA allows a much broader range of missions and architectures to be explored, but with less cost fidelity.





Study Assumptions: Satellite Measurements

Visible imaging of the Northern Hemispheres of all satellites at 1 km resolution, and at least one at 100 m resolution. We chose Miranda as the primary target for detailed study, with Titania or Oberon as secondary.

Visible/Near-IR spectroscopy of satellites with the above resolutions.

At least one close flyby of at least one satellite for gravity measurements and high resolution imaging.

We note that the visible and near-IR imagers desired for satellite science can also make important atmospheric measurements.




Shown without solar arrays

Architectures Explored in the RMA Study

- A: Minimum cost (Voyager-class) flyby.
- B: Upgraded flyby (includes 1- μ rad imager, VIMS-type instrument).
- C: Minimum cost flyby with a probe.
- D: Minimum cost flyby with 3 probes.
- E: Minimum cost flyby with 10 free-flying magnetometers.
- F: Minimum polar orbiter (Ka-band radio, simple magnetometer)
- G: New Frontiers orbiter (dual-band radio, enhanced magnetometer).
- H: Moderate orbiter (Option G with SWIR and microwave sounder).
- I: Cassini-class orbiter (instruments and range of orbit inclinations).
- J: Cassini-class orbiter with probe.
- K: Dual orbiter mission, one polar one equatorial.


Study Results: Cost vs. Science Value

Each architecture was judged against its ability to meet science goals related to the interior, atmosphere, magnetosphere, satellites, and rings.




Option "G" is the NF Orbiter, "H" is the Moderate Orbiter.

Study Results: Launch Masses for Chemical Propulsion




Option "G" is the NF Orbiter.

Study Results: Launch Masses for Solar Electric Propulsion



Option "G" is the NF Orbiter.

Study Results: Cost Estimates (Uncertainty +30%, -5%)




Option "G" is the NF Orbiter.

Study Results: Launch Year vs. Inserted Mass

For Solar-Electric Propulsion (SEP), all years have similar performance, and there is a trade off to be made among flight time (8-12 years), delivered mass (800 to 2000 kg), and launch vehicle.

For chemical propulsion, the launch-year is more important, with 2018 providing a particularly good geometry for a Jupiter gravity assist.



Study Results: General Conclusions

- Large masses can be placed into orbit around Uranus.
For example, using an Atlas 521 and only chemical propulsion, a **dry mass** in excess of 1500 kg (~100 kg for science instruments) can be inserted after a 12-year flight.

There are many trade-offs possible among cost, flight time, and delivered mass. Electric propulsion may be an attractive option.
- Uranus will be encountered near northern Solstice. This samples the same atmospheric season as Voyager did, but allows for imaging the unseen hemispheres of the satellites.
- Solar powered missions are feasible. Power is a significant constraint, but batteries, radioisotope heating units, and phasing of instrument on/off times allow the needed science return with solar panels producing only 100 W.
- Missions **may** be possible under the current New Frontiers (NF) cost cap.

Outline of Today's Talk


I) Introduction

II) The scientific importance of ice giants

III) General results from our study of Uranus missions


IV) Specifics of a Uranus Interior Mission

V) Conclusions



A Possible New Frontiers Mission (1 of 2)


- We found the most cost-effective, scientifically compelling mission to be an orbiter for high resolution mapping of the gravity and magnetic fields as a probe of interior structure.
- Our rough cost estimate ($\pm 30\%$), including all reserves, is 10% over the current NF guideline (\$650 million not counting the launch vehicle).
- Mass is not a limiting factor, so foreign contributions of instruments or a probe can be a way to increase science return while minimizing cost.
- Mission is possible with no new technology, though we need to optimize Ultraflex arrays for low light and temperatures.
- Advances in low-power electronics, improved downlink rates, low-temperature propellants, or aero-capture significantly improve capabilities.



A Possible New Frontiers Mission (2 of 2)

We found this mission scenario to be a good starting point for future studies:

- Launch September 2018 on an Atlas 521.
- Flybys of Venus (2), Earth, and Jupiter.
- Arrival at Uranus in September 2030.
- Insertion into a polar orbit ($\sim 70^\circ$ inclination).
- 1.2 year mission consisting of 10, 44-day elliptical orbits. Periapse 1.1 Uranus radii, apoapse 100 radii.



Science floor instrument package consists of


- X/Ka radio transmitters (Doppler tracking used for mapping the gravity field).
- Scalar and vector magnetometers (plus boom with star tracker).
- PEPPSI-type instrument for particle measurements.

Total science mass ~ 22 kg, not counting radio transmitters. 12 Gb of data generated during Uranus operations.

Subject to power, data volume, and cost constraints, ~ 100 kg of additional payload can be accommodated. An opportunity for satellite science!



Outline of Today's Talk

- I) Introduction
- II) The scientific importance of ice giants
- III) General results from our study of Uranus missions
- IV) Specifics of a Uranus Interior Mission
- V) Conclusions



What do we Ask of the Decadal Survey?

- Recommend that a mid-sized mission to Uranus or Neptune be launched in the next decade. This is crucial for understanding the diversity of planets (and satellites) and their formation and evolution.
- Conduct a more detailed mission study of the science capabilities and cost of a Uranus orbiter, including the option of using solar power.
- Assess our team's conclusion that a Uranus orbiter, focused on studies of the gravity and magnetic fields, is the most cost-effective, scientifically compelling mid-sized mission.




Hofstadter: Uranus Orbiter


Backup Slides




Types of Oceans in Icy Worlds



Europa:
*warm salty H₂O,
mantle contact*





Ganymede & Callisto:
perched salty H₂O(-NH₃?)



Titan: *open CH₄ seas*



Titan, Triton, large KBOs, and mid-sized icy satellites:
cold NH₃-H₂O, some perched, some mantle contact



Enceladus:
*cold H₂O-NH₃
or hydrothermal?*



Earth:
open salty H₂O

Study Assumptions: Mission Architecture

Do not use nuclear power sources (but allow radioactive heating units).

Keep approach velocities under 15 km/s (faster speeds make flyby encounters too brief, and orbiter missions difficult to slow down).


For hardware reliability, keep mission length under 15 years.

Use current technologies (e.g. aero-capture is not an option).


Consider launches between 2015 and 2023.

Orbital and flyby geometries must match the chosen instrument suite and mission objectives.

As seen from approaching spacecraft





As seen from (roughly) ecliptic north




For launches in the 2016 - 19 time frame and transfer times of 11-12 years, δ is $\sim 70^\circ$, decreasing with time such that for 2023 launches it is $< 60^\circ$. δ this high precludes encountering multiple satellites with a single flyby spacecraft.



Uranus: The Big Picture



Orton et al.



Hammel et al.
Icarus 2005



Weighting Functions and Atmospheric Profile

